

PHYSICAL STRUCTURE OF PLANETARY NEBULAE. III. THE LARGE AND EVOLVED NGC 1360

DANIEL B. GOLDMAN, MARTÍN A. GUERRERO¹, YOU-HUA CHU², AND ROBERT A. GRUENDL
 Astronomy Department, University of Illinois at Urbana-Champaign, 1002 West Green Street, Urbana, IL 61801
 dgoldman@uiuc.edu, mar@astro.uiuc.edu, chu@astro.uiuc.edu, gruendl@astro.uiuc.edu
The Astronomical Journal, 2004 November Issue

ABSTRACT

NGC 1360 is a large planetary nebula (PN) without an obvious shell morphology. We have analyzed H α images and high-dispersion echelle spectra of NGC 1360 in order to construct spatio-kinematic models and to determine its density distribution. The best-fit model indicates that NGC 1360 is a prolate ellipsoidal shell whose major axis is twice as long as its minor axis and is tilted by 60° with respect to the line of sight. The large kinematic age of the shell, $\sim 10,000$ yr, and the low density of the nebula, ≤ 130 H-atom cm⁻³, imply that NGC 1360 is an evolved PN and has begun to merge with the interstellar medium. The observed morphology and surface brightness profiles of NGC 1360 can be described well as a thick shell with a Gaussian radial density profile without a sharp inner edge, indicating a lack of on-going compression by a fast stellar wind. The FLIERs observed in NGC 1360 near the end of its major axis expand faster and are younger than the nebular shell.

Subject headings: ISM: kinematics and dynamics – ISM: planetary nebulae: general – planetary nebulae: individual (NGC 1360)

1. INTRODUCTION

Planetary nebulae (PNe) exhibit a wide range of morphologies. While the overall shape of a PN may be round, elliptical, or bipolar, most are limb-brightened, indicating a lower density in the interior, or the existence of a central cavity. It has been suggested that PNe are formed by dynamical interactions between the current fast stellar wind and previous slow wind (Kwok, Purton, & Fitzgerald 1978). Hydrodynamic calculations of interacting-stellar-wind models have indeed been successful in reproducing the hollow-shell structure and overall shapes of PNe by varying the pole-to-equator density gradient in the Asymptotic Giant Branch (AGB) progenitor’s slow wind (e.g., Frank & Mellema 1994). A shortcoming of the hydrodynamic modeling is that often several sets of parameters based on differing geometries are capable of reproducing the same morphology. Observations of spatially resolved kinematics of a PN can help differentiate among models.

We have been studying the physical structure of PNe using both imaging and high-dispersion spectroscopic observations, in order to understand their formation and evolution. Thus far, we have analyzed the Owl Nebula (NGC 3587) in Paper I (Guerrero et al. 2003) and NGC 7662 in Paper II (Guerrero et al. 2004). The Owl Nebula is a triple-shell PN; its “owl face” morphology is caused by bipolar cavities excavated by a fast wind that ceased some time ago, and its kinematics indicate that the nebular material is backfilling these cavities. NGC 7662 is also a triple-shell PN; its bright main nebula consists of two elliptical shells with simple structures in H α and [O III], although the outer rim of the main nebula is interspersed with [N II]-bright knots. These two PNe were chosen for our study because their morphologies have been

successfully reproduced by interacting-stellar-wind models.

There exist PNe that do not possess morphological features to suggest the presence of wind-wind interactions. NGC 1360 is such a PN. Its surface brightness does not dip deeply at the center or rise steeply at the limb to indicate a hollow shell structure. The lack of shell morphology in NGC 1360 makes it particularly interesting. Therefore, we have chosen this PN for a spatio-kinematic analysis in order to determine its physical structure and to investigate its evolutionary status. This paper reports our observations and analysis of NGC 1360.

2. OBSERVATIONS

2.1. Direct Images

Narrow-band H α images of NGC 1360 were obtained on 1999 October 8–9 with the 1 m telescope at Mount Laguna Observatory (MLO). The central wavelength and FWHM of the H α filter are 6563 Å and 17 Å, respectively. The detector was a Loral 2K CCD with a pixel size of 15 μ m, corresponding to an image scale of 0''.4 pixel⁻¹ and providing a field of view of 13'.3 $\times$ 13'.3. The angular resolution was $\sim 2''.8$, as determined from the FWHM of field stars in the images. A total of 10 images, each with an exposure time of 600 s, were combined to form the final image. This image is shown in Figure 1.

We have also retrieved archival *Hubble Space Telescope* images of NGC 1360 taken with the Wide Field and Planetary Camera 2 (WFPC2) and the F656N (H α) and F658N ([N II]) filters. These observations were obtained on 1994 February 15, during the “pre-cooldown” period when the WFPC2 was operating at a temperature of -77 °C (rather

¹ Current address: Instituto de Astrofísica de Andalucía, Consejo Superior de Investigaciones Científicas (CSIC), Apartado Correos 3004, E-18080, Granada, Spain.

² Visiting Astronomer, Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatories, operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under a cooperative agreement with the National Science Foundation.

than the optimal -88°C at which it normally operates). This higher operating temperature produced a substantially greater number of warm pixels and a higher dark current. To process the observations we first flagged and interpolated the warm pixels using the hot pixel lists available at STScI³, and then proceeded with the standard routines in the STSDAS package within IRAF. These images are not shown in this paper because the F658N image did not detect any nebulosity and the F656N image does not show any small-scale features unresolved in our ground-based $\text{H}\alpha$ image in Figure 1. The F656N image has only been used to flux-calibrate our ground-based $\text{H}\alpha$ image. The total $\text{H}\alpha$ flux from NGC 1360 is $\sim 3.1 \times 10^{-10}$ ergs cm^{-2} s^{-1} .

2.2. High-Dispersion Spectroscopic Observations

High-dispersion spectroscopic observations of NGC 1360 were made using the echelle spectrograph on the 4 m telescope at Cerro Tololo Inter-American Observatory (CTIO) on 2001 December 25. The 79 line mm^{-1} echelle grating and the long-focus red camera were used to achieve a reciprocal dispersion of $3.4 \text{ \AA mm}^{-1}$. The spectrograph was used in the single-order, long-slit mode to allow the coverage of the $\text{H}\alpha$ and $[\text{N II}] \lambda\lambda 6548, 6584$ lines along a slit length of $\sim 3'$. The data were recorded with the SITe 2K No. 6 CCD with a pixel size of $24 \mu\text{m}$, which corresponds to $0''.254 \text{ pixel}^{-1}$ along the spatial direction and $3.7 \text{ km s}^{-1} \text{ pixel}^{-1}$ along the dispersion direction. The slit width was $1''.0$, and the resultant instrumental FWHM was 12 km s^{-1} at the $\text{H}\alpha$ line.

The echelle observations were made along the major axis of NGC 1360 at the position angle (PA) of 27° . Three slit positions were used to cover the full extent of the major axis, as marked in Figure 1. The integration time at each slit position was 600 s. The angular resolution, determined by the seeing, was $\sim 2''.0$. A mosaic of the $\text{H}\alpha$ echellograms is shown in Figure 2, alongside an $\text{H}\alpha$ image oriented such that the major axis is parallel to the spatial direction of the echellogram.

3. RESULTS

3.1. Morphology

As shown in the $\text{H}\alpha$ image (Fig. 1), NGC 1360 appears elliptical with its major axis oriented along PA $\sim 30^\circ$. It lacks a sharp outer boundary because the surface brightness declines with distance until it drops below the background noise. In order to estimate the size of NGC 1360, we determined the background noise in the $\text{H}\alpha$ image and defined the nebular edge to be at 3σ above the background. Using the threshold of $3\sigma = 5.3 \times 10^{-16}$ ergs cm^{-2} s^{-1} arcsec^{-2} , NGC 1360 extends $9' \times 7'$, as shown by the contour in Figure 1. At a distance of $350_{-150}^{+1,000}$ pc as determined from the *Hipparcos* measurements (Acker et al. 1998), the linear size of NGC 1360 would be 0.9×0.7 pc.

While the overall surface brightness distribution of NGC 1360 falls off radially, there are depressions of $\sim 20\%$ within $30''$ of the central star and in the eastern hemisphere. The spatial distribution of these surface brightness depressions do not appear to be random. The central

depression extends into the eastern hemisphere and curves southward, while another distinct depression curves from the east to the north of the region surrounding the central star. The cause of these depressions is either lower gas densities or a higher extinction. If the latter is true, the non-random distribution of the depressions suggests that the obscuring material is internal to the nebula. A quantitative analysis of the surface brightness of NGC 1360 is presented in §4.

At the northern end of the major axis there exist several nebular knots extending from $240''$ out to $350''$ from the central star. Near the southern end of the major axis, there are also knots (marked by an arrow in Fig. 1), but they are fainter and detected at just the 3σ level. The faint southern knots, the central star, and the bright northern knots are not aligned along a straight line; instead, they form a slightly curved arc with the star at the apex. Such a configuration has been observed in other PNe and suggested nebular motion through the interstellar medium.

3.2. Kinematics

The echelle observations at the three consecutive slit positions have been mosaicked into an echellogram spanning the full major axis of NGC 1360 and are presented in Figure 2. This $\text{H}\alpha$ echellogram shows a position-velocity ellipse, indicating an expanding shell in the nebula. Furthermore, the position-velocity ellipse is tilted, with the southwestern and northeastern tips blue- and red-shifted, respectively. For an expanding ellipsoidal shell, the tilt of the position-velocity ellipse results from the inclination of the shell's polar axis with respect to the sky plane. As the position-velocity tilt occurs along the major axis of NGC 1360, its ellipsoidal shell must be prolate in shape, with its southwestern tip oriented toward us and its northeastern tip away from us.

The velocity profile of the $\text{H}\alpha$ line near the central star shows two broad components (see Fig. 3). The approaching component has a heliocentric velocity of $V_{\text{hel}} = 25 \text{ km s}^{-1}$ and a width of $\text{FWHM} = 30 \text{ km s}^{-1}$, while the receding component has $V_{\text{hel}} = 73 \text{ km s}^{-1}$ and $\text{FWHM} = 37 \text{ km s}^{-1}$. These widths are much larger than the thermal width of the $\text{H}\alpha$ line at 10^4 K , $\text{FWHM} = 21.4 \text{ km s}^{-1}$, indicating the existence of significant velocity variations along the line of sight within the nebular shell.

The average velocity of the approaching and receding sides of the nebula, $V_{\text{hel}} = 49 \text{ km s}^{-1}$, is adopted as the systemic velocity of NGC 1360. This result is in good agreement with the previous measurement of $V_{\text{hel}} = +49.1 \text{ km s}^{-1}$ by Bohuski & Smith (1974) using a Fabry-Perot scanner for a $60''$ aperture near the center of NGC 1360. The line-splitting at the center of a PN is usually adopted as twice its expansion velocity so that the apparent expansion velocity of NGC 1360 would be 24 km s^{-1} ; however, as we discuss in §4.1, this is an oversimplification of the expansion of NGC 1360.

In the $\text{H}\alpha$ line, both the nebula and the knots are detected; however, in the $[\text{N II}]$ line, only the knots are detected. This is illustrated by the $\text{H}\alpha$ and $[\text{N II}]$ lines of the northeastern end of NGC 1360's major axis shown in Figure 4. The low $[\text{N II}]/\text{H}\alpha$ ratio of the nebula is expected because the central star of NGC 1360 has an ef-

³ <http://www.stsci.edu/instruments/wfpc2/wfpc2.hotpix.html>

fective temperature of 110,000 K (Napiwotzki 1999) and the ionization stage of N in the nebula is higher than N^+ . The high $[N\ II]/H\alpha$ ratios of the knots suggest that they may be FLIERS, i.e., fast, low-ionization emission regions (Balick et al. 1993). Figure 4 shows that the innermost knots detected within the slit are at $\sim 260''$ from the central star and have a radial velocity of $V_{\text{hel}} \sim 100\text{ km s}^{-1}$ corresponding to a systemic velocity of $\sim +50\text{ km s}^{-1}$. The radial velocity of the knots increases outward by $0.5\text{ km s}^{-1}\text{ arcsec}^{-1}$. These knots clearly move faster than the shell expansion, confirming their identification as FLIERS.

4. DISCUSSION

4.1. Spatio-Kinematic Structure of NGC 1360

The $H\alpha$ image and echelle spectra of NGC 1360 can be used to construct spatio-kinematic models. In contrast to the identifiable thin shells in the Owl Nebula (Paper I) and NGC 7662 (Paper II) that can be modeled easily, the shell of NGC 1360 has both large velocity widths and irregular brightness variations in the echelle line image, which require simplifying assumptions to make the modeling tractable. We have assumed that NGC 1360 consists of a series of concentric thin ellipsoidal shells and the shell expansion velocity increases outwards. Consequently, the envelope of the position-velocity ellipse (marked in Fig. 2a) corresponds to an outer thin ellipsoidal shell that may be identified with the elliptical isophotes with a matching spatial extent in the direct image (marked in Fig. 2b). This outer thin shell can then be modeled using the methodology outlined by Guerrero et al. (2003), by assuming that the shell expands homologously (i.e., a Hubble law).

We model this shell of NGC 1360 as a prolate ellipsoidal shell with its polar axis tilted with respect to the line-of-sight by an inclination angle i . Its polar and equatorial radii are r_p and r_e , and its polar and equatorial expansion velocities are V_p and V_e , respectively. For the assumed homologous expansion, r_p/r_e is equal to V_p/V_e , and the kinematic age is simply r_e/V_e or r_p/V_p . A grid of models are constructed for a range of inclination angles i , axial ratios r_p/r_e , and equatorial expansion velocities V_e . These models are then used to synthesize nebular isophotes and to generate position-velocity ellipses for visual comparisons with the observations. Our best-fit model is marked in Figure 2, and has the following parameters: $i = 60^\circ$, $V_e = 35\text{ km s}^{-1}$ at $r_e = 130''$, $V_p = 70\text{ km s}^{-1}$ at $r_p = 260''$, and $t_{\text{kin}} = 10,000\text{ yr}$. Note that the inclination angle and the axial ratio are correlated so that lower values of i yield higher ratios of V_p/V_e . Despite this fact, the inclination angle and the axial ratio are both well constrained. The former is within $\pm 10^\circ$ from the optimal value, while the latter is within ± 0.25 from $V_p/V_e = 2.0$. The expansion velocities and kinematic age, on the other hand, are not as well constrained. For $i = 60^\circ$, the uncertainty in V_e is $\pm 10\text{ km s}^{-1}$, corresponding to about $\pm 30\%$.

Bear in mind that this model describes only an outer layer of the nebular shell of NGC 1360. The bulk of the nebular material is interior to this layer and expands more slowly. This densest shell material is responsible for the peaks in the velocity profile. For an inclination angle of 60° , the velocity split near the nebular center roughly corresponds to twice the equatorial expansion velocity of the densest nebular material in NGC 1360, 24 km s^{-1} (see

Fig. 3 and §3.2). However, the detailed geometry and kinematics of this dense material cannot be more precisely determined because the peaks of the line profiles do not form a well-defined position-velocity ellipse in the echelle image.

The FLIERS in NGC 1360 are projected along the apparent major axis of the nebula; thus, it is reasonable to assume that the FLIERS are distributed along the polar axis with an inclination angle of 60° . The apparent expansion velocities of the FLIERS, $\geq 50\text{ km s}^{-1}$, then correspond to deprojected expansion velocities of $\geq 100\text{ km s}^{-1}$. These expansion velocities are typical of those observed in FLIERS (Balick, Preston, & Icke 1987).

4.2. Physical Structure of NGC 1360

We can use the spatio-kinematic model of NGC 1360 deduced in §4.1 along with the observed $H\alpha$ surface brightness distribution to study the density structure within this nebula. The observed $H\alpha$ surface brightness profiles along several PAs are plotted in Figure 5. The depression in surface brightness seen in the eastern hemisphere of NGC 1360 is clearly visible in the profiles along PAs between 105° and 165° . In contrast, the surface brightness of the western hemisphere of NGC 1360 shows smoother profiles, without obvious “bumps” or “dents.” We have carried out the analysis of the density structure of NGC 1360 using the sections of the nebula between PA 225° and 345° because their uniformity best represents the overall distribution of material in the nebular shell.

The profiles of the western regions of NGC 1360 in Figure 5 exhibit a Gaussian shaped brightness distribution peaking at $40''$ – $60''$ and declining smoothly with increasing radius. The smooth shapes of the surface brightness profiles indicate that the nebula does not contain any regions with sharp density or extinction variations in the western hemisphere. These surface brightness distributions, in conjunction with the geometric information determined from the spatio-kinematic modeling in §3.2, can be used to derive the nebular density distribution. We have modeled the surface brightness profile assuming a Gaussian density distribution and fitted this model to the observed $H\alpha$ surface brightness profiles between PA 225° and 345° . The best-fit model peaks at $35''$ with a FWHM of $60''$ along the equatorial plane and peaks at $70''$ with a FWHM of $100''$ along the polar direction. The best-fit models are overplotted in Figure 5: good agreements are obtained from PA 225° to 345° . Using the flux calibrated surface brightness we further determined that the peak density is $\sim 130\text{ H-atom cm}^{-3}$, and that the density at $r_e = 130''$ is $\sim 0.1\text{ H-atom cm}^{-3}$. The low density at the outer edge of the nebula implies that it is beginning to merge into the interstellar medium. The total mass in the model, $\sim 0.4\text{ M}_\odot$, is estimated assuming that the surface brightness depressions in the eastern hemisphere are caused by internal dust absorption. If the lower surface brightness is caused by a lack of material in the shell, the mass would be over-estimated, but this error is small as compared to the uncertainty in mass due to the large error bar in the distance to NGC 1360.

5. SUMMARY AND CONCLUSIONS

We have obtained images and high-dispersion long-slit spectra of NGC 1360 in order to study its spatio-kinematic structure. We have modeled an outer layer of the shell as a prolate ellipsoid and find a major-to-minor axial ratio of 2:1 and a polar axis tilted by 60° with respect to the line of sight. The kinematic age derived from this model is $\sim 10,000$ yr.

We have found that the observed morphology and surface brightness profiles of NGC 1360 can be modeled well by a thick shell with a Gaussian radial density profile. This smooth density distribution implies that there is no sharp inner edge as expected from compression by a fast stellar wind. If a fast wind was present in the past and compressed the AGB wind into a thin shell, that shell has since diffused. The low nebular density, ≤ 130 H-atom

cm^{-3} , suggests that NGC 1360 is an evolved PN; its low-density (0.1 H-atom cm^{-3}) envelope has begun to merge with the interstellar medium.

The FLIERs observed in NGC 1360 are intriguing. The kinematic age of the FLIERs, based on their de-projected distances from the central star, ≥ 0.5 pc, and their de-projected expansion velocities, $\geq 100 \text{ km s}^{-1}$, is $\sim 5,000$ yr. This kinematic age is about $1/2$ that of the PN itself. NGC 1360 is one of the very few cases of evolved PNe in which collimated outflows are observed (e.g., Corradi et al. 1999).

We thank Jin-chung Hsu and Stefano Casertano for their help in addressing the warm pixel problem of the *HST* WFPC2 image.

REFERENCES

- Acker, A., Fresneau, A., Pottasch, S. R., & Jasiewicz, G. 1998, A&A, 337, 253
 Balick, B., Preston, H. L., & Icke, V. 1987, AJ, 94, 1641
 Balick, B., Rugers, M., Terzian, Y., & Chengalur, J. N. 1993, ApJ, 411, 778
 Bohuski, T. J. & Smith, M. G. 1974, ApJ, 193, 197
 Corradi, R. L. M., Perinotto, M., Villaver, E., Mampaso, A., & Gonçalves, D. R. 1999, ApJ, 523, 721
 Frank, A. & Mellema, G. 1994, ApJ, 430, 800
 Guerrero, M. A., Chu, Y.-H., Manchado, A., & Kwitter, K. B. 2003, AJ, 125, 3213 (Paper I)
 Guerrero, M. A., Jaxon, E. G., & Chu, Y.-H. 2004, AJ, in press, October issue (Paper II)
 Kwok, S., Purton, C. R., & Fitzgerald, P. M. 1978, ApJ, 219, L125
 Napiwotzki, R. 1999, A&A, 350, 101

FIG. 1.— An $H\alpha$ image of NGC 1360 displayed with two different intensity contrasts to highlight the faint outer regions (*left*) and the structure in the central regions (*right*). The contour in the left panel corresponds to the 3σ level above the background. The slit positions used in the echelle spectroscopic observations are marked on the image. The faint southern FLIERS are marked with arrows in both panels.

FIG. 2.— (*left*) $H\alpha$ echellogram along the major axis of NGC 1360. The geocoronal $H\alpha$ line is marked. (*right*) $H\alpha$ image of NGC 1360 oriented to match its major axis with the echelle slit position for direct comparison. The jagged contours overlaid on the echellogram and direct image have matching spatial extent and correspond to isophotes detected at 4σ and 7σ levels, respectively. The smooth ellipses overlaid on the echellogram and direct image are based on our best-fit spatio-kinematic model. The bright FLIERS northeast to the nebula are visible in both panels.

FIG. 3.— Velocity profile of the $H\alpha$ line within the central region of NGC 1360. The position of the geocoronal $H\alpha$ line is marked. The systemic velocity adopted for NGC 1360 is also marked.

FIG. 4.— An expanded view of the $H\alpha$ and $[N\ II]$ echellograms at the northeast end of NGC 1360 to show the FLIERS. The spatial position is measured relative to the central star. The $H\alpha$ line shows both the nebular shell and the FLIERS, while the $[N\ II]$ line shows only the FLIERS. The FLIERS exhibit a velocity gradient along the slit visible in both lines.

FIG. 5.— Surface brightness profiles (solid lines) of the $H\alpha$ line extracted at 30° intervals. The peak at the zero position corresponds to the central star, and the position angle is marked for the profile on either side of the star. The dashed curves are the profiles from our best-fit model.

This figure "fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0407568v1>

This figure "fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0407568v1>

This figure "fig3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0407568v1>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0407568v1>

This figure "fig5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0407568v1>